

R-matrices for the semicyclic representations of $U_q\widehat{sl}(2)$

I.T.Ivanov* and D.B.Uglov

Department of Physics, State University of New York at Stony Brook
Stony Brook, NY 11794-3800

9 March 1992

Abstract

R-matrices for the semicyclic representations of $U_q\widehat{sl}(2)$ are found as a specialization in the R-matrices of the checkerboard chiral Potts model

The chiral Potts model [1][2] is the first example of a new class of integrable statistical mechanical models whose Boltzmann weights are meromorphic functions on an algebraic variety of genus greater than 1. From the mathematical point of view this model is related to the theory of cyclic representations of the quantum algebra $U_q\widehat{sl}(2)$ at $q^N = 1$ [3]. This relation is expressed by the fact that the R-matrix intertwining two tensor products of cyclic representations is built up of the Boltzmann weights of the chiral Potts model [3][4].

In this paper we consider certain limit of the chiral Potts model which in the algebraic language corresponds to the descent from the cyclic representation to the semicyclic one [6]. Boltzmann weights remain well defined in this limit and as a result we obtain R-matrices which intertwine semicyclic representations of $U_q\widehat{sl}(2)$ and satisfy the Yang-Baxter equation.

Series of R-matrices were obtained for semicyclic representations of $U_qsl(2)$ in [7][8]. The R-matrices obtained here for the larger algebra $U_q\widehat{sl}(2)$ may be specialized to $U_qsl(2)$ subalgebra. We checked that in the case $N = 3$ the R-matrix from [7] is equivalent by a similarity transform to the R-matrix obtained as a specialization in the chiral Potts model in this letter.

*e-mail iti@dirac.physics.sunysb.edu

1. Checkerboard chiral Potts R-matrix The chiral Potts model Boltzmann weights

$$\frac{W_{\xi,\eta}(i)}{W_{\xi,\eta}(i-1)} = \frac{\mu_\xi y_\eta - x_\xi \omega^i}{\mu_\eta y_\xi - x_\eta \omega^i}, \quad \frac{\overline{W}_{\xi,\eta}(i)}{\overline{W}_{\xi,\eta}(i-1)} = \mu_\xi \mu_\eta \frac{x_\xi \omega - x_\eta \omega^i}{y_\eta - y_\xi \omega^i} \quad (1)$$

depend on two points ξ and η of two identical algebraic curves C_ξ and C_η defined by the relations

$$\mu^N = \frac{k'}{1 - kx^N} = \frac{1 - ky^N}{k'} \quad (2)$$

and the same for C_η . Also $k^2 + k'^2 = 1$.

We shall associate [4] with every cyclic representation π_ξ of the quantum algebra $U_q \widehat{sl}(2)$ two copies of the curve (2) which we shall denote by C_ξ and $C_{\xi'}$. The variables of the sets $\xi = x, y, \mu$ and $\xi' = x', y', \mu'$ as well as k are functions of the parameters of the representation. Then the R-matrix intertwining tensor products of two cyclic representations π_ξ and π_η is given by [4] :

$$R(\xi, \eta) = \sum_{i,j,k,l=1}^N \omega^{i(j-k)} (Z^j \otimes Z^k) (X^{-1} \otimes X)^{i+l} \times \overline{W}_{\xi,\eta}(k) \overline{W}_{\xi',\eta'}(j) \widehat{W}_{\xi',\eta}(i) \widehat{W}_{\xi,\eta'}(l) \quad (3)$$

where Z and X are N by N matrices satisfying :

$$ZX = \omega XZ \quad Z^N = X^N = 1 \quad \omega = q^2 \quad (4)$$

and $\widehat{W}_{\xi,\eta}$ is a Fourier transform of the Boltzmann weight (1) : $\widehat{W}_{\xi,\eta}(i) = \sum_{k=1}^N W_{\xi,\eta}(k) \omega^{-ik}$.

$$\frac{\widehat{W}_{\xi,\eta}(i)}{\widehat{W}_{\xi,\eta}(i-1)} = \frac{\mu_\eta y_\xi - \mu_\xi y_\eta \omega^{i-1}}{\mu_\eta x_\eta - \mu_\xi x_\xi \omega^i} \quad (5)$$

Throughout this paper we consider only the case N odd .

Matrix elements of the R-matrix (3) are vertex weights of the checkerboard chiral Potts model [2], [5]

2. Taking limit in the Boltzmann weights. Here we take a special limit in the checkerboard chiral Potts model which will allow us to get the R-matrices for the semicyclic representations. It is shown in the appendix A that taking

the semicyclic limit in the cyclic representation is equivalent to the following rescaling of the variables :

$$k = \varepsilon^{-N} K, \quad k' = \varepsilon^{-N} K' \quad (6)$$

$$x = \varepsilon X, \quad y = \varepsilon^{-1} Y, \quad \mu = \varepsilon^{-1} M, \quad x' = \varepsilon^{-1} X', \quad y' = \varepsilon Y', \quad \mu' = \varepsilon M' \quad (7)$$

and subsequently setting ε to zero. The Boltzmann weights entering the R-matrix are invariant under this rescaling :

$$\frac{W_{\xi', \eta}(i)}{W_{\xi', \eta}(i-1)} = \frac{M_{\xi'} Y_{\eta} - X_{\xi'} \omega^i}{M_{\eta} Y_{\xi'} - X_{\eta} \omega^i} \quad \frac{W_{\xi, \eta'}(i)}{W_{\xi, \eta'}(i-1)} = \frac{M_{\xi} Y_{\eta'} - X_{\xi} \omega^i}{M_{\eta'} Y_{\xi} - X_{\eta'} \omega^i} \quad (8)$$

$$\frac{\overline{W}_{\xi, \eta}(i)}{\overline{W}_{\xi, \eta}(i-1)} = M_{\xi} M_{\eta} \frac{X_{\xi} \omega - X_{\eta} \omega^i}{Y_{\eta} - Y_{\xi} \omega^i} \quad \frac{\overline{W}_{\xi', \eta'}(i)}{\overline{W}_{\xi', \eta'}(i-1)} = M_{\xi'} M_{\eta'} \frac{X_{\xi'} \omega - X_{\eta'} \omega^i}{Y_{\eta'} - Y_{\xi'} \omega^i} \quad (9)$$

The finite parts of the parameters $X, Y \dots$ obey the following equations :

$$M^N = \frac{k'}{1 - K X^N} = \frac{\varepsilon^{2N} - K Y^N}{K'} \quad M'^N = \frac{k'}{\varepsilon^{2N} - K X'^N} = \frac{1 - K Y'^N}{K'} \quad (10)$$

with $K^2 + K'^2 = \varepsilon^{2N}$. Let us now take the limit $\varepsilon \rightarrow 0$. The Boltzmann weights (7-8) remain well defined. The equations of the curve show that M is proportional to Y and M' is proportional to $1/X'$. So the Boltzmann weights (8,9) simplify and to describe them more neatly we introduce one more set of parameters A, B, C, D defined by

$$A = K^{1/N} Y^{-1} \quad B = -K^{1/N} X \quad C = K^{1/N} X'^{-1} \quad D = -K^{1/N} Y' \quad (11)$$

This set of parameters satisfy the equations of two Fermat curves :

$$A^N = B^N + 1 \quad C^N = D^N + 1 \quad (12)$$

The Boltzmann weights entering the R-matrix finally have the following form :

$$\frac{W_{\xi, \eta'}(i)}{W_{\xi, \eta'}(i-1)} = \frac{D_{\eta} - B_{\xi} \omega^i}{C_{\eta} - A_{\xi} \omega^i} \quad \frac{W_{\xi', \eta}(i)}{W_{\xi', \eta}(i-1)} = \frac{C_{\xi} - A_{\eta} \omega^i}{D_{\xi} - B_{\eta} \omega^i} \quad (13)$$

$$\frac{\overline{W}_{\xi, \eta}(i)}{\overline{W}_{\xi, \eta}(i-1)} = \frac{B_{\xi} \omega - B_{\eta} \omega^i}{A_{\xi} - A_{\eta} \omega^i} \quad \frac{\overline{W}_{\xi', \eta'}(i)}{\overline{W}_{\xi', \eta'}(i-1)} = \frac{C_{\eta} \omega - C_{\xi} \omega^i}{D_{\eta} - D_{\xi} \omega^i} \quad (14)$$

The periodicity property $W(i+N) = W(i)$, $\overline{W}(i+N) = \overline{W}(i)$ is guaranteed by the equations (12).

3. R-matrix for the semicyclic representations of $U_q\widehat{sl}(2)$ Here we present the R-matrix and associated Hamiltonian.

The R-matrix intertwining two semicyclic representations π_ξ and π_η (See Appendix A) as a result of the previous section is also given by the formula (3) with Boltzmann weights (13,14). The intertwining condition can be proved along the same lines as in the periodic case [4]. To prove the Yang-Baxter equation we introduce four auxiliary Boltzmann weights :

$$\frac{W_{\xi,\eta}(i)}{W_{\xi,\eta}(i-1)} = \frac{1 - \phi B_\xi A_\eta \omega^i}{1 - \phi A_\xi B_\eta \omega^i} \quad \frac{W_{\xi',\eta'}(i)}{W_{\xi',\eta'}(i-1)} = \frac{\phi C_\xi D_\eta - \omega^i}{\phi D_\xi C_\eta - \omega^i} \quad (15)$$

$$\frac{\overline{W}_{\xi',\eta}(i)}{\overline{W}_{\xi',\eta}(i-1)} = \frac{\omega - \phi C_\xi B_\eta \omega^i}{1 - \phi D_\xi A_\eta \omega^i} \quad \frac{\overline{W}_{\xi,\eta'}(i)}{\overline{W}_{\xi,\eta'}(i-1)} = \frac{\phi B_\xi C_\eta \omega - \omega^i}{\phi A_\xi D_\eta - \omega^i} \quad (16)$$

We checked that the Boltzmann weights (13-14),(15-16) satisfy a set of eight star-triangle equations

$$\sum_{d=1}^N \overline{W}_{q,r}(b-d) W_{p,r}(a-d) \overline{W}_{p,q}(d-c) = R_{p,q,r} W_{p,q}(a-b) \overline{W}_{p,r}(b-c) W_{q,r}(a-c) \quad (17)$$

with p, q and r replaced by both primed and unprimed indices and ϕ arbitrary number. This set of equations imply the Yang-Baxter equation for the semicyclic R-matrix of the form (3). The fact that the R-matrix (3) satisfies the Yang-Baxter equation can also be derived from the intertwining property and an indecomposability of the third-order tensor product of the semicyclic representations of $U_q\widehat{sl}(2)$ at generic values of the parameters. If the restrictions $D = -qA, B = -q^{-1}C$ are imposed on the parameters A, B, C, D we obtain $e_0 = f_1$, and $f_0 = e_1$ in the representation of $U_q\widehat{sl}(2)$ (see Appendix A). And the resulting R-matrix intertwines between the representations of finite dimensional quantized algebra $U_qsl(2)$. In the case $N = 3$ we checked that this R-matrix is equivalent to the R-matrix of [7] by the similarity transformation.

The spin-chain hamiltonian corresponding to the R-matrix (3) can be obtained using the standard procedure [2]. To write down it we introduce the following operators:

$$X_i = \overset{1}{1} \otimes \cdots \otimes \overset{i}{X} \otimes \cdots \otimes \overset{L}{1} \quad Z_i = \overset{1}{1} \otimes \cdots \otimes \overset{i}{Z} \otimes \cdots \otimes \overset{L}{1} \quad (18)$$

where $i = 1, \dots, L$; L is the length of the periodic lattice, X and Z are the same as in (4). In terms of these operators the hamiltonian reads as follows:

$$H = \sum_{j=1}^L h_{j,j+1}, \quad (19)$$

$$h_{j,j+1} = \sum_{n=1}^N F(n) X_j^{-n} X_{j+1}^n + \sum_{n,m=1}^N [G_+(n,m) X_j^{-n} X_{j+1}^n Z_j^m + G_-(n,m) X_j^{-n} X_{j+1}^n Z_{j+1}^m] \quad (20)$$

where the explicit form of the coefficients F, G_+, G_- is given in the appendix B. The form of this hamiltonian is the same as of the general *checkerboard* chiral Potts model. In the case of the well-known chiral Potts model the coefficients $G_+(n, m)$ and $G_-(n, m)$ become proportional to $\delta_{n,0}$ and the hamiltonian acquires the usual form [1].

Appendix A

In this appendix we describe the $U_q \widehat{sl}(2)$ algebra and its cyclic and semicyclic representations at $q^N = 1$. After that we describe in detail the limiting procedure used in section 2.

Quantized enveloping algebra $U_q \widehat{sl}(2)$ is generated by the elements t_i, t_i^{-1}, e_i and f_i ($i = 0, 1$) subject to the relations

$$t_i t_i^{-1} = t_i^{-1} t_i = 1 \quad t_i t_j = t_j t_i \quad (21)$$

$$t_i e_j t_i^{-1} = q^{a_{ij}} e_j, \quad t_i f_j t_i^{-1} = q^{-a_{ij}} f_j, \quad [e_i, f_j] = \frac{t_i - t_i^{-1}}{q_i - q_i^{-1}} \delta_{ij} \quad (22)$$

with $a_{ij} = 2, i = j$ and $a_{ij} = -2, i \neq j$. These generators also obey the Serre relations.

Tensor products of representations are formed using the following comultiplication

$$\Delta(e_0) = e_0 \otimes 1 + z^{-1} t_0 \otimes e_0, \quad \Delta(e_1) = e_1 \otimes 1 + z t_1 \otimes e_1 \quad (23)$$

$$\Delta(f_0) = f_0 \otimes t_0^{-1} + z \otimes f_0, \quad \Delta(f_1) = f_1 \otimes t_1^{-1} + z^{-1} \otimes f_1 \quad (24)$$

$$\Delta(t_i) = t_i \otimes t_i, \quad \Delta(z) = z \otimes z \quad (25)$$

z is central element commuting with the algebra $U_q \widehat{sl}(2)$. N -dimensional cyclic representation of the quantum algebra $U_q \widehat{sl}(2)$ may be described in terms of the two matrices Z and X and five complex parameters $\xi = (a, b, x_0, x_1, c)$,

$$\pi_\xi(e_0) = x_0 X^{-1} \frac{b^2 Z^{-1} - 1}{q - q^{-1}}, \quad \pi_\xi(f_0) = (abx_0)^{-1} \frac{a^2 Z - 1}{q - q^{-1}} X \quad (26)$$

$$\pi_\xi(f_1) = (abx_1)^{-1} X^{-1} \frac{b^2 Z^{-1} - 1}{q - q^{-1}}, \quad \pi_\xi(e_1) = x_1 \frac{a^2 Z - 1}{q - q^{-1}} X \quad (27)$$

$$\pi_\xi(t_0) = \frac{bZ^{-1}}{qa}, \quad \pi_\xi(t_1) = \frac{qaZ}{b}, \quad \pi_\xi(z) = c \quad (28)$$

R-matrix is an operator which intertwins tensor products of two representations π_ξ and π_η taken in an alternated order :

$$R(\xi, \eta)(\pi_\xi \otimes \pi_\eta)\Delta(g) = (\pi_\eta \otimes \pi_\xi)\Delta(g)R(\xi, \eta), \text{ for all } g \in U_q\widehat{sl}(2). \quad (29)$$

Necessary and sufficient condition for an intertwiner between tensor products of two cyclic representations π_ξ and π_η to exist is that the following four quantities should be the same in both representations.

$$\gamma_1 = \frac{x_1^N(a^{2N} - 1)}{1 - a^N b^{-N} c^N}, \quad \gamma_2 = \frac{a^{2N} - 1}{a^N b^N x_0^N (a^N b^{-N} - c^N)} \quad (30)$$

$$\gamma_3 = \frac{b^{2N} - 1}{a^N b^N x_1^N (a^{-N} b^N - c^{-N})}, \quad \gamma_4 = \frac{x_0^N (b^{2N} - 1)}{1 - a^{-N} b^N c^{-N}} \quad (31)$$

These four invariants are not independent but obey $\gamma_1 \gamma_3 = \gamma_2 \gamma_4$. In order to write the R-matrix for the cyclic case in the factorized form a change of parameters $\{\gamma_1, \gamma_2, \gamma_3, \gamma_4, a, b, c, x_0, x_1\} \mapsto \{k, k_0, k_1, x, y, \mu, x', y', \mu'\}$ is convenient.

$$k^2 = \frac{-1}{\gamma_1 \gamma_3}, \quad k_0^N = -k \gamma_4, \quad k_1^N = -k \gamma_1 \quad (32)$$

$$x_0 = k_0 x', \quad x_1 = k_1 y', \quad a^2 = \frac{x \mu \mu'}{y'}, \quad b^2 = \frac{y}{x' \mu \mu'}, \quad c = ab \frac{y'}{qx} \quad (33)$$

Then the R-matrix satisfying (29) is given by (3) with the chiral Potts Boltzmann weights.

We will call N-dimensional representation of $U_q\widehat{sl}(2)$ semicyclic if e_1^N and f_0^N are represented by non-zero scalars while $e_0^N = f_1^N = 0$. In terms of the parameters of the cyclic representation (26-28) this means $b^{2N} - 1 = 0$. Let $\pi_\xi \xi = (a, b, c, x_0, x_1)$ and $\pi_\eta \eta = (\hat{a}, \hat{b}, \hat{c}, \hat{x}_0, \hat{x}_1)$ are two cyclic representations. Let also

$$b^{2N} - 1 = \varepsilon^{2N} f, \quad \hat{b}^{2N} - 1 = \varepsilon^{2N} \hat{f} \quad (34)$$

where f and $\hat{f}$ are some finite functions of the parameters of the corresponding representations and ε is a number. In limit the $\varepsilon \rightarrow 0$ both representations π_ξ and π_η become semicyclic. Let us using the equations (32-33) to explicitly single out powers of ε from the invariants and representation parameters :

$$\gamma_3 = \varepsilon^{2N} \Gamma_3 = \varepsilon^{2N} f G_3, \quad \gamma_4 = \varepsilon^{2N} \Gamma_4 = \varepsilon^{2N} f G_4, \quad \gamma_1 = \Gamma_1, \quad \gamma_2 = \Gamma_2 \quad (35)$$

$$k = \varepsilon^{-N} K, \quad k_0 = \varepsilon K_0, \quad k_1 = \varepsilon^{-1} K_1 \quad (36)$$

$$x = \varepsilon X, \quad y = \varepsilon^{-1} Y, \quad \mu = \varepsilon^{-1} M, \quad x' = \varepsilon^{-1} X', \quad y' = \varepsilon Y', \quad \mu' = \varepsilon M' \quad (37)$$

As is mentioned in the main text the Boltzmann weights are invariant under this rescaling and we may safely set ε to zero. The invariants Γ_1 and Γ_2 still play the role of necessary conditions for existing of the intertwiner. The initial invariant conditions for γ_3 and γ_4 become an defining equation for the functions f and $\hat{f}$ from (34):

$$\frac{f}{\hat{f}} = \frac{\hat{G}_3}{G_3} = \frac{\hat{G}_4}{G_4} \quad (38)$$

The Boltzmann weights entering R-matrix (3) do not depend on the particular choice of f and $\hat{f}$ satisfying (38). The four other possible Boltzmann weights become trivial in the semicyclic limit :

$$\frac{W_{\xi,\eta}(i)}{W_{\xi,\eta}(i-1)} = 1 \quad \frac{W_{\xi',\eta'}(i)}{W_{\xi',\eta'}(i-1)} = 1 \quad \frac{\overline{W}_{\xi',\eta}(i)}{\overline{W}_{\xi',\eta}(i-1)} = \omega \quad \frac{\overline{W}_{\xi,\eta'}(i)}{\overline{W}_{\xi,\eta'}(i-1)} = 1 \quad (39)$$

Appendix B

Here we list the coefficients G_+, G_-, F entering the hamiltonian (20):

$$F(n) = \sum_{i=1}^N \widehat{W}_{\xi,\xi'}(n-i) \widehat{W}_{\xi',\xi}(i) \left[\sum_{s=1}^{n-i} \frac{\gamma - \delta \omega^{s-1}}{C_\xi - D_\xi \omega^s} - \sum_{s=1}^i \frac{\beta - \alpha \omega^{s-1}}{B_\xi - A_\xi \omega^s} \right]; \quad (40)$$

$$G_+(n, m) = E(n, m) \overline{W}_{\xi',\xi}(m); \quad G_-(n, m) = E(n, -m) \overline{W}_{\xi,\xi}(m) \quad (41)$$

where

$$E(n, m) = \sum_{i=1}^N \omega^{im} \widehat{W}_{\xi,\xi'}(i) \widehat{W}_{\xi',\xi}(n-i); \quad (42)$$

$$(43)$$

and

$$\overline{W}_{\xi',\xi'}(i) = \frac{\gamma}{C_\xi} \frac{(\frac{C_\xi}{D_\xi})^i}{1 - \omega^i}; \quad \overline{W}_{\xi,\xi}(i) = \frac{\beta}{B_\xi} \frac{(\frac{B_\xi}{A_\xi})^i}{1 - \omega^{-i}}; \quad (44)$$

$$\widehat{W}_{\xi',\xi}(i) = \prod_{s=1}^i \frac{C_\xi - D_\xi \omega^{s-1}}{A_\xi - B_\xi \omega^s}; \quad \widehat{W}_{\xi,\xi'}(i) = \prod_{s=1}^i \frac{D_\xi - C_\xi \omega^{s-1}}{B_\xi - A_\xi \omega^s}; \quad (45)$$

where the parameters $A_\xi, B_\xi, C_\xi, D_\xi, \alpha, \beta, \gamma, \delta$ satisfy the following equations:

$$A_\xi^N = B_\xi^N + 1; \quad C_\xi^N = D_\xi^N + 1; \quad (46)$$

$$\alpha = \left(\frac{B_\xi}{A_\xi}\right)^{N-1} \beta; \quad \gamma = \left(\frac{D_\xi}{C_\xi}\right)^{N-1} \delta; \quad (47)$$

Summary. Quantum enveloping algebras at roots of unity have hierarchy of irreducible multiparameter representations : cyclic $\rightarrow$ semicyclic $\rightarrow$ nilpotent representations. In this letter we have shown that the process of specializing the parameters in the cyclic representation of $U_q\widehat{sl}(2)$ to make it semicyclic goes through on the R-matrix level also. We believe that the R-matrices from [8] are equivalent to the R-matrices obtained here when the later are specialized to the $U_qsl(2)$ case. The limit to the nilpotent representations [10] is under investigation.

The R-matrices for the cyclic representations of $U_q\widehat{sl}(2)$ are already known [9] and it seems feasible to find the semicyclic limit in this case.

Acknowledgement

We are grateful to prof. B.M.McCoy for many helpful suggestions and to R.Kedem for discussions.

References

- [1] H.Au-Yang, B.M.McCoy, J.H.H.Perk, S.Tang and M.Yan, Phys. Lett. A 123 (1987) 219; B.M.McCoy, J.H.H.Perk, S.Tang and C.H.Sah, Phys.Lett. A 125 (1987) 9
- [2] R.J.Baxter, J.H.H.Perk and H.Au-Yang, Phys.Lett. A 128 (1988) 138
- [3] V.V.Bazhanov and Y.G.Stroganov, J.Stat.Phys. 59 (1990) 799
- [4] E.Date, M.Jimbo, M.Miki and T.Miwa, RIMS preprint 706 (1990)
- [5] R.J.Baxter, V.V.Bazhanov and J.H.H.Perk, Int.J.Mod.Phys.B 4 (1990) 803
- [6] P.Roche and D.Arnaudon, Lett. Math. Phys. 17 (1989) 295; G.Keller, Lett. Math. Phys. 21 (1991) 273
- [7] C.Gomez, M.Ruiz-Altaba and G.Sierra, Phys.Lett.B 265 (1991) 95
- [8] C.Gomez and G.Sierra, Nucl.Phys.B (to appear)
- [9] E.Date, M.Jimbo, K.Miki and T.Miwa, RIMS preprint 715 (1990)
- [10] A. Berkovich, C. Gomez and G. Sierra, preprint 1992